

Percent of strains susceptible:

Gram Negative Enteric Bacteria	No. Tested	Ampicillin/Sulbactam	Piperacillin/tazobactam	Aztreonam	Cefazolin	Cefoxitin	Ceftriaxone	Cefepime	Meropenem	Gentamicin	Tobramycin	Amikacin	Levofloxacin	Sulfa/Trimethoprim	Nitrofurantoin	Tetracycline
<i>Enterobacter cloacae</i>	64				R	R		97	98	100	100	100	97	95	39	94
<i>Escherichia coli</i>	284	64	96	90	87	89	90	90	100	88	91	100	81	76	95	78
<i>Klebsiella pneumoniae</i>	94	65	90	67	66	88	67	67	97	96	94	100	93	66	R	74
<i>Proteus mirabilis</i>	68	93	100	94	90	96	94	94	100	96	97	100	74	75	R	R

Gram Negative Non-enteric Bacteria	No. Tested	Piperacillin/tazobactam	Cetazidime	Cefepime	Meropenem	Aztreonam	Gentamicin	Tobramycin	Amikacin	Levofloxacin	Minocycline	Sulfa/Trimethoprim
<i>Pseudomonas aeruginosa</i>	200	91	80	80	90	92	91	97	98	77		
<i>Stenotrophomonas maltophilia</i>	37		31							68	100	92

Enterococci/Streptococci	No. Tested	Penicillin/ampicillin	Levofloxacin	Vancomycin	Linezolid	Nitrofurantoin	Gentamicin Synergy	Streptomycin Synergy
<i>Enterococcus faecalis</i>	185	100	83	96	100	100	78	82
<i>Enterococcus faecium</i>	43	R	R	28	100	R	97	63
<i>Streptococcus agalactiae</i>	86	100	97	100	100			

Gram Positive Bacteria	No. Tested	Penicillin/ampicillin	Oxacillin/nafcillin	Erythromycin	Clindamycin	Gentamicin*	Levofloxacin	Sulfa/Trimethoprim	Vancomycin	Rifampin*	Tetracycline	Minocycline	Linezolid	Nitrofurantoin
<i>Staphylococcus aureus</i>	483	R	51	37	86	100	65	93	100	99	92	100	100	99
<i>Staphylococcus epidermidis</i>	53	R	R	R	94	79	47	38	100	98	77	74	100	96

*Rifampin and gentamicin should be used only in combination with another agent

R denotes intrinsic resistance

Dosing recommendations are generalized; dosing should be optimized for each patient.