

Gram Negative Enteric Bacteria	No. Tested	Ampicillin/Sulbactam	Piperacillin/tazobactam	Aztreonam	Cefazolin	Cefoxitin	Ceftriaxone	Cefepime	Meropenem	Gentamicin	Tobramycin	Amikacin	Levofloxacin	Sulfa/Trimethoprim	Nitrofurantoin	Tetracycline
<i>Escherichia coli</i>	738	67	98	97	95	96	97	97	100	94	94	100	89	81	99	84
<i>Klebsiella pneumoniae</i>	91	92	98	95	95	98	95	95	100	100	100	100	98	89	33	87
<i>Proteus mirabilis</i>	35	91	100	100	97	100	100	100	100	91	91	100	80	83	R	R

Gram Positive Bacteria	Number Tested	Penicillin/ampicillin	Oxacillin/nafcillin	Erythromycin	Clindamycin	Gentamicin*	Levofloxacin	Sulfa/Trimethoprim	Vancomycin	Rifampin*	Tetracycline	Minocycline	Linezolid	Nitrofurantoin
<i>Staphylococcus aureus</i>	120	R	52	43	95	100	74	98	100	100	97	100	100	100
<i>Streptococcus agalactiae</i>	35	100		34	86		97		100		R		100	

*Rifampin and gentamicin should be used only in combination with another agent

Enterococci	No. Tested	Penicillin/ampicillin	Levofloxacin	Vancomycin	Linezolid	Nitrofurantoin	Gentamicin Synergy	Streptomycin Synergy
<i>Enterococcus faecalis</i>	79	96	87	99	100	100	85	89

R denotes intrinsic resistance

Dosing recommendations are generalized; dosing should be optimized for each patient.

Adam Warner, PharmD
Antimicrobial Stewardship Pharmacist
(269) 341-6230
warnera@bronsonhg.org

Glenda Pietryga, PharmD
Pharmacy Manager
(269) 657-1481
PIETRYGG@bronsonhg.org

Richard A. Van Enk, Ph.D.
Epidemiologist
(269) 341-6316
vanenkr@bronsonhg.org