

Percent of strains susceptible:

Gram Negative Enteric Bacteria	No. Tested	Ampicillin/Sulbactam	Piperacillin/tazobactam	Aztreonam	Cefazolin	Cefoxitin	Ceftriaxone	Cefepime	Meropenem	Gentamicin	Tobramycin	Amikacin	Levofloxacin	Sulfa/Trimethoprim	Nitrofurantoin	Tetracycline
<i>Escherichia coli</i>	694	62	97	96	95	95	96	96	100	95	95	100	91	80	98	81
<i>Klebsiella pneumoniae</i>	88	88	98	98	98	95	98	98	99	100	100	100	100	95	R	R
<i>Proteus mirabilis</i>	31	97	100	100	97	100	100	100	100	97	97	100	87	94	R	R

Gram Positive Bacteria	Number Tested	Penicillin/ampicillin	Oxacillin/nafcillin	Erythromycin	Clindamycin	Gentamicin	Levofloxacin	Sulfa/Trimethoprim	Vancomycin	Rifampin*	Tetracycline	Minocycline	Linezolid	Nitrofurantoin
<i>Staphylococcus aureus</i>	118	R	46	33	94	100	69	97	100	100	97	100	100	100
<i>Streptococcus agalactiae</i>	42	100	R	R	74	R	98	R	100	R	R	R	100	R

\*Rifampin should be used only in combination with another agent

Enterococci	No. Tested	Penicillin/ampicillin	Levofloxacin	Vancomycin	Linezolid	Nitrofurantoin	Gentamicin Synergy	Streptomycin Synergy
<i>Enterococcus faecalis</i>	80	100	88	99	100	99	86	86

R denotes intrinsic resistance

Dosing recommendations are generalized; dosing should be optimized for each patient.

Adam Warner, PharmD  
 Antimicrobial Stewardship Pharmacist  
 (269) 341-6230  
[warnera@bronsonhg.org](mailto:warnera@bronsonhg.org)

Glenda Pietryga, PharmD  
 Pharmacy Manager  
 (269) 657-1481  
[PIETRYGG@bronsonhg.org](mailto:PIETRYGG@bronsonhg.org)

Richard A. Van Enk, Ph.D.  
 Epidemiologist  
 (269) 341-6316  
[vanenkr@bronsonhg.org](mailto:vanenkr@bronsonhg.org)